

Indications for Removable Partial Dentures: A Literature Review

Bernd Wöstmann, Dr Med Dent^a/Ejvind Budtz-Jørgensen, DDS, Dr Odont^b/Nick Jepson, BDS, PhD, FDS^c/
Eiko Mushimoto, DDS, Dr Med Dent, PhD^d/Sigvard Palmqvist, LDS, Odont Dr/PhD^e/
Afrodite Sofou, DDS, Dr Dent^f/Bengt Öwall, LDS, Odont Dr, Dr Med HC/PhD^g

Purpose: The purpose of this report was to evaluate indications for the use of removable partial dentures based on a comprehensive literature review. **Materials and Methods:** Using a model similar to a Delphi process, the literature relating to the indications and contraindications for the prescription of removable partial dentures was discussed by seven experienced educators in a 2.5-day workshop. **Results:** Evidence for indications and contraindications for the prescription of removable partial dentures is not clearly stated in the literature; however, some basic principles are defined. There appears to be a trend in favor of the use of the shortened dental arch concept or implant-supported restorations instead of conventional removable partial dentures, given the evidence that the long-term use of removable dentures is associated with increased risks of caries and periodontitis and low patient acceptance. The presence of sound abutment teeth appears to encourage the use of removable partial dentures, as the fixed partial denture alternative requires sacrifice of healthy hard tissues. When economic factors influence the decision-making process, removable partial dentures are often chosen. **Conclusion:** While evidence-based indications and contraindications for prescribing removable partial dentures are still lacking, major underlying principles for clinical decision making are identified. *Int J Prosthodont* 2005;18:139–145.

The context of increasing life spans and evidence from various national dental health surveys in industrialized countries indicate that the proportion of edentulous people will continue to decline and that more people will retain more teeth into old age.^{1,2} Projections based on data from such surveys^{2,3} suggest

a decline in tooth loss but an increased need for management of partial edentulism in patients with compromised oral or general health status. This trend of age-related increased tooth retention suggests that partially edentulous cohorts will be older than before and probably less disposed than younger people to such treatment with extensive tooth- or implant-supported fixed partial dentures (FPD). Consequently, socioeconomic factors and population trends suggest increased future treatment needs with different partial prostheses.

The present review is based on the hypothesis that the decision to prescribe a removable partial denture (RPD) is mainly based on a subjective mix of professional, functional, cultural, and economic considerations rather than on compelling scientific evidence. Therefore, the authors sought to identify evidence-based indications for RPDs.

Materials and Methods

The seven authors who evaluated the literature are clinical academic prosthodontists with at least 18 years' active practice experience.

^aProfessor, Department of Prosthodontics, Justus Liebig University, Giessen, Germany.

^bProfessor Emeritus, Division of Gerodontology and Removable Prosthodontics, Section of Dental Medicine, University of Geneva, Switzerland.

^cSenior Lecturer, Restorative Dentistry, School of Dental Sciences, University of Newcastle upon Tyne, United Kingdom.

^dAssociate Professor, Department of Prosthodontics, Iwate Medical University College of Dentistry, Japan.

^eAssociate Professor, Department of Prosthetic Dentistry, Faculty of Health Sciences, University of Copenhagen, Denmark.

^fAssociate Professor, Department of Prosthetic Dentistry, School of Dentistry, Aristotle University, Thessaloniki, Greece.

^gProfessor and Chair, Department of Prosthetic Dentistry, Faculty of Health Sciences, University of Copenhagen, Denmark.

Correspondence to: Dr Bernd Wöstmann, Justus Liebig University, Dental Clinic, Schlängenzahl 14, Giessen D-35392, Germany. Fax: + 49 641 99 46139. e-mail: Bernd.Woestmann@dentist.med.uni-giessen.de

Table 1 Key Words Included in Electronic Search

Key words	Hits	Key words	Hits
RPD	283	Satisfaction AND RPD	15
		Removable denture	168
Nutrition AND dental	3,523	Nutrition AND denture	170
Subsearch: AND (tooth or teeth) AND NOT child*	468		
Satisfaction AND dental	2,365	Satisfaction AND denture	763
Subsearch: AND prosth*	521		
Quality of life AND dental	695	Quality of life AND denture	113
QoL AND dental	32	QoL AND denture	10

Search Strategy

An online search was conducted using the key words listed in Table 1 in the following databases:

- Cochrane Database of Abstracts of Cochrane Reviews
- MEDLINE
- PubMed
- Current Contents
- DIMDI (German Institute for Documentation in Medicine)

The search was augmented by a manual search of relevant journals and textbooks unavailable through online databases. Case reports and technical procedure descriptions were excluded from the analysis. From the remaining references, those that focused on the following areas related to RPDs were selected:

- Masticatory function
- Nutritional status
- Quality of life and patient satisfaction
- Interactions between RPDs and oral structures (caries, periodontitis, residual ridge resorption)
- Temporomandibular disorders (TMD)
- Prophylactic aspects (TMD, tooth wear, tooth migration)
- Alternative treatment options
- Contraindications

Literature Analysis

The literature analysis was organized in several rounds in a manner similar to the Delphi principle:

- Round 1: literature search according to the strategy described
- Round 2: identification of the literature to be included in the evaluation in a workshop held in Giessen, Germany, in May 2002

- Round 3: circulation of a preliminary draft paper and inclusion of additional new published literature
- Round 4: circulation of the draft and inclusion of additional literature
- Round 5: approval of the final paper

The databases were first searched in April 2002; additional searches followed during rounds 4 and 5 (May and October 2003).

Results

Masticatory Function

The key determinants of masticatory performance are the number of functional tooth units and the bite force (Table 2). An age-related decrease of masticatory performance is associated with tooth loss, a decline of muscle mass and density of the masticatory muscles, and frailty.⁴⁻⁶ However, patients' subjective assessment of chewing is influenced more by age than dental and prosthodontic status.⁷

Masticatory performance improves after prosthodontic treatment independent of the design and type (FPD/RPD) of the denture.^{8,9} However, unless the patient has fewer than three occluding pairs of posterior teeth, there seems to be no sociofunctional benefit to be gained from replacing missing posterior teeth.¹⁰

Nutritional Status

Nutritional deficiency is common in older adults.¹¹ The impaired masticatory function associated with tooth loss may lead to a deficient and unbalanced diet,^{12,13} malnutrition, and even compromised general health. However, other factors, such as systemic health, socioeconomic status, and dietary habits, will also influence dietary selection (Table 2). Although masticatory performance generally increases after prosthodontic treatment, food selection remains unchanged irrespective of the prosthesis design.

Table 2 Evidence-Based Statements Drawn from Literature Review

Statement	Supporting studies	Cumulated N*
Masticatory function		
Masticatory performance is determined by number of occlusal units and bite force	38-40	644
Chewing performance is affected by tooth loss and muscle status	4-6,39-42	11,305
Masticatory performance improves after prosthodontic treatment	8,9,43-46	568
Nutritional status		
Impaired masticatory function compromises general health	11,12,30,47-49	3,617
Impaired masticatory function influences food selection	50,51	735
Socioeconomic factors and general health influence food selection	52,53	> 166
Prosthodontic treatment does not influence dietary intake	8,39,54-56	793
Quality of life and patient satisfaction		
RPD patients are more dissatisfied with their oral conditions than those with FPDs or natural teeth	16,18,19,57-59	8,100
Many RPDs are not used	22,60-63	629
RPDs cause inconvenience	22,61,64,65	321
Interactions between RPDs and oral structures		
There is an association between the use of RPDs and caries and periodontal disease	33,66-75	8,137
If plaque control is achieved in appropriately designed RPDs, the risks for caries and periodontitis are not increased	36,76-80	3,394
There is an association between ill-fitting RPDs and bone resorption	22,81	227
TMD		
Absence of molar support is not an etiologic factor for TMD	20-22,66,82,83	1,419
Prophylactic aspects		
Unrestored shortened dental arches are occlusally stable	84,85	851

*Total sample size of referenced studies where available; > = total sample size not available.

RPD = removable partial denture; FPD = fixed partial denture; TMD = temporomandibular disorders.

While a correlation between dental and nutritional status is evident, it remains unclear what the dependent and independent variables are. This observation is aggravated by patients' tendency to overestimate their nutritional status and masticatory performance.¹⁴ Most patients consider themselves to be adequately nourished and tend to assess their dentures more favorably than a qualified prosthodontist would.¹⁵

Quality of Life and Patient Satisfaction

A significant proportion of prescribed RPDs are not used. In this context, it must be clearly recognized that practical problems with RPDs (food retention, pressure spots, etc) are common and may be the reason so many patients stop wearing their RPDs (Table 2). Patient satisfaction increases when the RPD adds a significant number of occlusal units to the dentition.¹⁷ However, the improved perceived function gained by an RPD replacing only a few teeth does not compensate for the discomfort of wearing the RPD. There is a reported superiority of FPDs with respect to patient satisfaction, but this does not in all aspects clearly favor FPDs over RPDs.^{18,19}

Interactions Between RPDs and Oral Structures

The use of RPDs is associated with an increased risk of caries and periodontitis, although there is no evidence

that RPDs per se cause damage. With appropriate design and adequate plaque control, long-term clinical service RPDs have no detrimental effects on the periodontium or abutment teeth. Caries, however, can be a particular problem in elderly persons. There is strong evidence of an association between the use of RPDs and root surface caries (Table 2).

TMD

Some studies suggest a correlation between missing molars and TMD²⁰; however, there is almost no evidence that TMD symptoms are associated with a shortened dental arch (SDA).²¹ Consequently, the replacement of missing posterior teeth by an RPD cannot be claimed to prevent the development of TMD (Table 2). In addition, poorly maintained RPDs may be associated with parafunctional jaw movements in the long term.²²

Prophylactic Aspects

A continuously used, well-fitting RPD can be expected to stabilize the occlusion and prevent tooth movement. However, occlusal stability in dentitions with unrestored SDAs is much higher than what has been traditionally assumed (Table 2). In this regard, no significant differences between SDA and RPD patients have been reported.

Alternative Treatment Options

Implant-supported prostheses are nearly always an alternative treatment choice for partially edentulous cases; therefore, this option has not been included in the present analysis. When only a few teeth are missing and the patient has no functional complaint, an SDA concept should be considered.²³ If the abutment teeth need crowning, attachment-retained (eg, precision attachment or telescopic) RPDs or cantilever FPDs may be suggested for better longevity and oral comfort.²⁴ If the teeth are sound, an acrylic resin-bonded FPD or resin-bonded adhesive attachment retaining the RPD is also an option.^{25,26} Whenever possible, fixed restorations should be considered as a first choice, as they demonstrate better survival.

Contraindications

Definitive contraindications to the use of RPDs have not been reported. Nevertheless, diseases such as diabetes and dry mouth may restrict RPD treatment because of the inability of the mucosa to withstand mechanical trauma. Esthetic considerations and aspects of comfort may also be contraindications.^{19,27} There is only weak evidence that an allergy to specific metals in the casting alloy also contraindicates an RPD.²⁸

Discussion

There appears to be a clear connection between oral and dental health status and the well-being of a patient, and numerous studies suggest that nutritional inadequacies may be related to dental status.^{29,30} However, the quality of supportive scientific evidence is not compelling. Nevertheless, it seems prudent to assert that adequate dental and prosthodontic status is an important contributory variable to a person's general health and—especially in industrialized countries—well-being and is significantly linked to socioeconomic status.³¹ Adequate prosthodontic rehabilitation clearly appears to increase chewing ability^{8,9} and enhance an edentulous patient's ability to select an appropriate diet; however, supporting evidence for a similar effect in the partially dentate is weak.

Given the evidence that oral comfort is more effectively achieved using fixed rather than removable restorations, an RPD is not the treatment of choice in an SDA when esthetics is not effected.³² The use of an RPD to replace only a few teeth where the improvement in oral function and/or esthetics is only marginal must therefore be questioned.³² The beneficial effects of an RPD on patient satisfaction are counteracted by the oral discomfort inevitably induced by the prosthesis itself, especially when only a small number of teeth

are missing. As the number of teeth to be replaced increases, the indication for the use of an RPD becomes stronger.

Treatment with RPDs is noninvasive to the remaining dentition's structure, but RPDs can increase the risk of caries, periodontitis, and residual ridge reduction, particularly in distal and extension areas. The specific risk of root caries when RPDs are used in elderly persons where the risk for inadequate oral hygiene is high and regular professional maintenance programs may be difficult suggests that RPD prescriptions should be avoided. In such patients, an SDA approach is a well-precedented treatment option.³³

In the management of patients with a diagnosed TMD, a distinction should be made between those reporting problems and pain and others without any complaints. If in the former group restoration of molar support is advisable, an RPD is the preferred treatment option given its advantage of reversibility.

A patient's financial status and third-party health care funding systems appear to strongly influence the choice of different prosthodontic treatment options, a rather difficult comparison at an international level. In general, inadequate fiscal resources generally favor the RPD, but other economic aspects have to be considered as well: In highly industrialized countries, an RPD is often regarded as a low-cost treatment option; however, in many other countries, even the simplest dental restoration may prove unaffordable. It is clear that systems of health care provision strongly influence the indications not only for RPDs, but for all types of restorations. Nevertheless, a correctly designed "hygienic" RPD³⁴ offers a comparatively low cost and noninvasive alternative to other highly sophisticated treatment options, although maintenance costs will increase in the long term to reflect both a shorter longevity of RPDs^{35,36} and the potential increased long-term treatment need. It is probably unlikely that the additional necessary expense will exceed the total initial and maintenance costs of more sophisticated restorations.

The pragmatic choice between different treatment options is based on both professional and subjective evaluations. Additional considerations include specifics regarding the burden of illness present, time for treatment, operator skills, patient perceptions, expected longevity, and, of course, expense. However, objective evaluations of most of these aspects are not available.³⁷

Conclusions

A detailed analysis of the literature failed to provide highly evidence-based indications for RPD treatment. Nevertheless, a lower level of evidence provided the following guidelines:

- Application of an SDA concept tends to preclude the indication for an RPD.
- The RPD is associated with an increased risk of caries and, to a lesser extent, periodontal disease, especially in the long term and in patients with poor oral health maintenance.
- Prophylactic oral health treatment with an RPD is questionable.
- The presence of sound abutment teeth in a well-maintained mouth strengthens the indication for a RPD.
- If provision of a conventional FPD is not possible because of extensive loss of teeth and implant treatment is not realistic, the clasp-retained RPD is a valid alternative.
- The application of optimal biologically related design principles may reduce RPD-related risks.
- The risk of low patient acceptance of an RPD must be considered.
- Economic considerations are strong indications for RPD treatment.

References

1. Kelly M, Steele J, Nuttal N, et al. Adult Dental Health Survey—Oral Health in the United Kingdom 1998. London: National Statistics Public Inquiry Service, 2000.
2. I+G Gesundheitsforschung. Bedarfsermittlung für prothetische Leistungen bis zum Jahr 2020. Munich, Germany: I+G Gesundheitsforschung, 2001.
3. Douglass CW, Watson AJ. Future needs for fixed and removable partial dentures in the United States. *J Prosthet Dent* 2002;87:9–14.
4. Miura H, Watanabe S, Isogai E, Miura K. Comparison of maximum bite force and dentate status between healthy and frail elderly persons. *J Oral Rehabil* 2001;28:592–595.
5. Tzakis MG, Österberg T, Carlsson GE. A study of some masticatory functions in 90-year old subjects. *Gerodontology* 1994;11:25–29.
6. Carlsson GE. Masticatory efficiency: The effect of age, the loss of teeth and prosthetic rehabilitation. *Int Dent J* 1984;34:93–97.
7. Lappalainen R, Nyssonen V. Self-assessed chewing ability of Finnish adults with removable dentures. *Gerodontology* 1987;3:238–241.
8. Kapur KK, Garrett NR, Dent RJ, Hasse AL. A randomized clinical trial of two basic removable partial denture designs. Part II: Comparisons of masticatory scores. *J Prosthet Dent* 1997;78:15–21.
9. Kapur KK. Veterans Administration Cooperative Dental Implant Study—Comparisons between fixed partial dentures supported by blade-vent implants and removable partial dentures. Part III: Comparisons of masticatory scores between two treatment modalities. *J Prosthet Dent* 1991;65:272–283.
10. Locker D, Clarke M, Payne B. Self-perceived oral health status, psychological well-being, and life satisfaction in an older adult population. *J Dent Res* 2000;79:970–975.
11. Appollonio I, Carabellese C, Fratola A, Trabucchi M. Influence of dental status on dietary intake and survival in community-dwelling elderly subjects. *Age Ageing* 1997;26:445–456.
12. Mojon P, Budtz-Jørgensen E, Rapin CH. Relationship between oral health and nutrition in very old people. *Age Ageing* 1999;28:463–468.
13. Österberg T, Steen B. Relationship between dental state and dietary intake in 70-year-old males and females in Göteborg, Sweden: A population study. *J Oral Rehabil* 1982;9:509–521.
14. Wickop H, Wöstmann B, Ferger P, Kolb G. Ernährungszustand und zahnärztlich-prothetische Versorgung älterer Patienten. *Swiss Dent* 1998;19:5–7.
15. Zlataric DK, Celebic A. Treatment outcomes with removable partial dentures: A comparison between patients and prosthodontic assessments. *Int J Prosthodont* 2001;14:423–426.
16. Kerschbaum T. Langzeitergebnisse und Konsequenzen. In: Hupfaut L (ed). *Praxis der Zahnheilkunde*, ed 6—Teilprothesen. Munich, Germany: Urban & Schwarzenberg, 1988.
17. van Waas M, Meeuwissen J, Meuwissen R, Käyser A, Kalk W, van't Hof M. Relationship between wearing a removable partial denture and satisfaction in the elderly. *Community Dent Oral Epidemiol* 1994;22:315–318.
18. Kapur KK. Veterans Administration Cooperative Dental Implant Study—Comparisons between fixed partial dentures supported by blade-vent implants and removable partial dentures. Part II: Comparisons of success rates and periodontal health between two treatment modalities. *J Prosthet Dent* 1989;62:685–703.
19. Jepson N, Allen F, Moynihan P, Kelly P, Thomason M. Patient satisfaction following restoration of shortened mandibular dental arches in a randomized controlled trial. *Int J Prosthodont* 2003;16:409–414.
20. Tallents RH, Macher DJ, Kyrkanides S, Katzberg RW, Moss ME. Prevalence of missing posterior teeth and intraarticular temporomandibular disorders. *J Prosthet Dent* 2002;87:45–50.
21. Türp JC, Strub JR. Prosthetic rehabilitation in patients with temporomandibular disorders. *J Prosthet Dent* 1996;76:418–423.
22. Witter DJ, De Haan AF, Käyser AF, van Rossum GM. A 6-year follow-up study of oral function in shortened dental arches. Part II: Craniomandibular dysfunction and oral comfort. *J Oral Rehabil* 1994;21:353–366.
23. van Waas MA, Meeuwissen JH, Meeuwissen R, Käyser AF. Oral function in dentate elderly with reduced dentitions. *Gerodontology* 1993;10:40–43.
24. Vermeulen AH, Keltjens HM, van't Hof MA, Käyser AF. Ten-year evaluation of removable partial dentures: Survival rates based on retreatment, not wearing and replacement. *J Prosthet Dent* 1996;76:267–272.
25. Watson RM, Gilmour AG. Modern solutions for limited tooth loss in the dental arch. *Eur J Prosthodont Restorative Dent* 1994;2:171–177.
26. Kern M. Care of young female patient using adhesive attachments and removable partial dentures [in German]. *Die Quintessenz* 1991;42:577–590.
27. Budtz-Jørgensen E, Bochet G, Grundman M, Borgis S. Aesthetic considerations for the treatment of partially edentulous patients with removable dentures. *Pract Periodontics Aesthet Dent* 2000;12:765–772.
28. Savion Y, Sharon-Buller A, Kalisker Y, Kalisker N, Sela M. The use of Dental D (polyacetal resin) as an alternative for chrome-cobalt removable partial denture: A case report [abstract]. *J Israel Dent Assoc* 1920;18:30–31, 108.
29. Nakata M. Masticatory function and its effects on general health. *Int Dent J* 1998;48:540–548.
30. Albiin N, Asplund K, Bjermer L. Nutritional status of medical patients on emergency admission to hospital. *Acta Med Scand* 1982;212:151–156.
31. Mericske-Stern R, Kowalski J. Nachsorgebefugung und Recallverhalten von älteren Patienten mit abnehmbaren Prothesen. *Schweiz Monatsschr Zahnmed* 1990;100:1053–1105.
32. Knezovic ZD, Celebic A, Valentic-Peruzovic M, Jerolimov V, Panduric J. A survey of treatment outcomes with removable partial dentures. *J Oral Rehabil* 2003;30:847–854.
33. Jepson NJ, Moynihan PJ, Kelly PJ, Watson GW, Thomason JM. Caries incidence following restoration of shortened lower dental arches in a randomized controlled trial. *Br Dent J* 2001;191:140–144.
34. Öwall B, Budtz-Jørgensen E, Davenport J, et al. Removable partial denture design: A need to focus on hygienic principles? *Int J Prosthodont* 2002;15:371–378.

35. Kerschbaum T, Leempoel PJB. Kronen und Brücken–Konsequenzen aus Langzeitergebnissen. In: Voß R, Meiners H (eds). *Fortschritte der Zahnärztlichen Prothetik und Werkstoffkunde*, ed 4. Munich, Germany: Hanser, 1989.
36. Wöstmann B. Tragedauer von Einstückgußprothesen im überwachten Gebrauch. *Dtsch Zahnärztl Z* 1997;52:100–104.
37. Strassburger C, Heydecke G, Kerschbaum T. Influence of prosthetic and implant therapy on satisfaction and quality of life: A systematic literature review. Part 1—Characteristics of the studies. *Int J Prosthodont* 2004;17:83–93.
38. Hatch JP, Shinkai RS, Sakai S, Rugh JD, Paunovich ED. Determinants of masticatory performance in dentate adults. *Arch Oral Biol* 2001;46:641–648.
39. Demers M, Bourdages J, Brodeur JM, Benigeri M. Indicators of masticatory performance among elderly complete denture wearers. *J Prosthet Dent* 1996;75:188–193.
40. Helkimo E, Carlsson GE, Helkimo M. Chewing efficiency and state of dentition. A methodologic study. *Acta Odontol Scand* 1978;36:33–41.
41. Witter DJ, Cramwinckel AB, van Rossum GM, Käyser AF. Shortened dental arches and masticatory ability. *J Dent* 1990;18:185–189.
42. Neill DJ, Phillips HI. The masticatory performance, dental state, and dietary intake of a group of elderly army pensioners. *Br Dent J* 1970;128:581–585.
43. Feine JS, Maskawi K, de Grandmont P, Donohue WB, Tanguay R, Lund JP. Within-subject comparisons of implant-supported mandibular prostheses: Evaluation of masticatory function. *J Dent Res* 1994;73:1646–1656.
44. van der Bilt A, Olthoff LW, Bosman F, Oosterhaven SP. Chewing performance before and after rehabilitation of post-canine teeth in man. *J Dent Res* 1994;73:1677–1683.
45. Nagasawa T, Tsuru H. A comparative evaluation of masticatory efficiency of fixed and removable restorations replacing mandibular first molars. *J Prosthet Dent* 1973;30:263–273.
46. Liedberg B, Spiechowicz E, Öwall B. Mastication with and without removable partial dentures: An intraindividual study. *Dysphagia* 1995;10:107–112.
47. Hildebrandt GH, Loesche WJ, Lin CF, Bretz WA. Comparison of the number and type of dental functional units in geriatric populations with diverse medical backgrounds. *J Prosthet Dent* 1995;73:253–261.
48. Loesche WJ, Abrams J, Terpenning MS, et al. Dental findings in geriatric populations with diverse medical backgrounds. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 1995;80:43–54.
49. Wayler AH, Muench ME, Kapur KK, Chauncey HH. Masticatory performance and food acceptability in persons with removable partial dentures, full dentures and intact natural dentition. *J Gerontol* 1984;39:284–289.
50. Sheiham A, Steele JG, Marcenes W, et al. The relationship among dental status, nutrient intake, and nutritional status in older people. *J Dent Res* 2001;80:408–413.
51. Elmståhl S, Birkhed D, Christiansson U, Steen B. Intake of energy and nutrients before and after dental treatment in geriatric long-stay patients. *Gerodontology* 1988;4:6–12.
52. Budtz-Jørgensen E, Chung JP, Mojón P. Successful aging—The case for prosthetic therapy. *J Public Health Dent* 2000;60:308–312.
53. Keller HH. Malnutrition in institutionalized elderly: How and why? *J Am Geriatr Soc* 1993;41:1212–1218.
54. Gunne HS. The effect of removable partial dentures on mastication and dietary intake. *Acta Odontol Scand* 1985;43:269–278.
55. Sebring NG, Guckes AD, Li SH, McCarthy GR. Nutritional adequacy of reported intake of edentulous subjects treated with new conventional or implant-supported mandibular dentures. *J Prosthet Dent* 1995;74:358–363.
56. Garrett NR, Kapur KK, Hasse AL, Dent RJ. Veterans Administration Cooperative Dental Implant Study—Comparisons between fixed partial dentures supported by blade-vent implants and removable partial dentures. Part V: Comparisons of pretreatment and post-treatment dietary intakes. *J Prosthet Dent* 1997;77:153–161.
57. Redford M, Drury TF, Kingman A, Brown LJ. Denture use and the technical quality of dental prostheses among persons 18–74 years of age: United States, 1988–1991. *J Dent Res* 1996;75:714–725.
58. Kapur KK. Veterans Administration Cooperative Dental Implant Study—Comparisons between fixed partial dentures supported by blade-vent implants and removable partial dentures. Part IV: Comparisons of patient satisfaction between two treatment modalities. *J Prosthet Dent* 1991;66:517–530.
59. Meeuwissen JH, van Waas MA, Meeuwissen R, Käyser AF, van't Hof MA, Kalk W. Satisfaction with reduced dentitions in elderly people. *J Oral Rehabil* 1995;22:397–401.
60. Kuboki T, Okamoto S, Suzuki H, et al. Quality of life assessment of bone-anchored fixed partial denture patients with unilateral mandibular distal-extension edentulism. *J Prosthet Dent* 1999;82:182–187.
61. Carlsson GE, Hedegård B, Koivumaa KK. Studies in partial dental prostheses. IV. Final results of a 4-year longitudinal investigation of dentogingivally supported partial dentures. *Acta Odontol Scand* 1965;23:443–472.
62. Jepson N, Allen F, Moynihan P, Kelly P, Thomason M. Patient satisfaction following restoration of shortened mandibular dental arches in a randomized controlled trial. *Int J Prosthodont* 2003;16:409–414.
63. Jepson NJ, Thomason JM, Steele JG. The influence of denture design on patient acceptance of partial dentures. *Br Dent J* 1995;178:296–300.
64. Peterhans G, Brunner T. Nachuntersuchung von Gerüstprothesen bei minderbemittelten Patienten—I. Resultate der Patientenbefragung. *Schweiz Monatsschr Zahnmed* 1986;96:482–499.
65. Peterhans G, Brunner T. Nachuntersuchung von Gerüstprothesen bei minderbemittelten Patienten—II. Resultate der Befunderhebung. *Schweiz Monatsschr Zahnmed* 1986;96:755–773.
66. Budtz-Jørgensen E, Isidor F. A 5-year longitudinal study of cantilevered fixed partial dentures compared with removable partial dentures in a geriatric population. *J Prosthet Dent* 1990;64:42–47.
67. Zlataric DK, Celebic A, Valentic-Peruzovic M. The effect of removable partial dentures on periodontal health of abutment and non-abutment teeth. *J Periodontol* 2002;73:137–144.
68. Chandler JA, Brudvik JS. Clinical evaluation of patients eight to nine years after placement of removable partial dentures. *J Prosthet Dent* 1984;51:736–743.
69. Lappalainen R, Koskenranta-Wuorinen P, Markkanen H. Periodontal and cariological status in relation to different combinations of removable dentures in elderly men. *Gerodontology* 1987;3:122–124.
70. Tuominen R, Ranta K, Paunio I. Wearing of removable partial dentures in relation to dental caries. *J Oral Rehabil* 1988;15:515–520.
71. Wright PS, Hellyer PH, Beighton D, Heath R, Lynch E. Relationship of removable partial denture use to root caries in an older population. *Int J Prosthodont* 1992;5:39–46.
72. Drake CW, Beck JD. The oral status of elderly removable partial denture wearers. *J Oral Rehabil* 1993;20:53–60.
73. Locker D, Ford J, Leake JL. Incidence of and risk factors for tooth loss in a population of older Canadians. *J Dent Res* 1996;75:783–789.
74. Steele JG, Walls AW, Murray JJ. Partial dentures as an independent indicator of root caries risk in a group of older adults. *Gerodontology* 1997;14:67–74.
75. Aquilino SA, Shugars DA, Bader JD, White BA. Ten-year survival rates of teeth adjacent to treated and untreated posterior bounded edentulous spaces. *J Prosthet Dent* 2001;85:455–460.
76. Petridis H, Hempton TJ. Periodontal considerations in removable partial denture treatment: A review of the literature. *Int J Prosthodont* 2001;14:164–172.

77. Kapur KK, Deupree R, Dent RJ, Hasse AL. A randomized clinical trial of two basic removable partial denture designs. Part I: Comparisons of five-year success rates and periodontal health. *J Prosthet Dent* 1994;72:268–282.
78. Bergman B, Ericson G. Cross-sectional study of the periodontal status of removable partial denture patients. *J Prosthet Dent* 1989;61:208–211.
79. Bergman B, Hugoson A, Olsson CO. A 25 year longitudinal study of patients treated with removable partial dentures. *J Oral Rehabil* 1995;22:595–599.
80. Vanzeveren C, D'Hoore W, Bercy P. Influence of removable partial denture on periodontal indices and microbiological status. *J Oral Rehabil* 2002;29:232–239.
81. Plotnick IJ, Beresin VE, Simkins AB. The effects of variations in the opposing dentition on changes in the partially edentulous mandible. Part I. Bone changes observed in serial radiographs. *J Prosthet Dent* 1975;33:278–286.
82. Budtz-Jørgensen E, Luan W, Holm-Pedersen P, Fejerskov O. Mandibular dysfunction related to dental, occlusal and prosthetic conditions in a selected elderly population. *Gerodontology* 1985;1:28–33.
83. Battistuzzi P, Käyser A, Kanters N. Partial edentulism, prosthetic treatment and oral function in a Dutch population. *J Oral Rehabil* 1987;14:549–555.
84. Witter DJ, De Haan AF, Käyser AF, van Rossum GM. A 6-year follow-up study of oral function in shortened dental arches. Part I: Occlusal stability. *J Oral Rehabil* 1994;21:113–125.
85. Sarita PT, Kreulen CM, Witter DJ, van't Hof M, Creugers NH. A study on occlusal stability in shortened dental arches. *Int J Prosthodont* 2003;16:375–380.

Literature Abstract

An in vitro evaluation of the reliability and validity of an electronic pantograph by testing with five different articulators

There have been a lot of efforts to transform the actual jaw movements on the articulators to recreate a patient's occlusion. The electronic mandibular tracing device may make the clinical procedure both simple and convenient. The purpose of this project was to evaluate the reliability and validity of the Cadiax Compact in calculating condylar settings for five different articulators: Denar D5A, Denar Mark II, Whip Mix 8500, Hanau Modular, and Panadent PCH. The Cadiax Compact sensors and styli were mounted to each articulator with custom-made mounting device that included a C-shaped aluminum plate, a round brass cross bar, and two adjustable side arms. At 3-, 5-, and 10-mm condylotrack distance, the condylar settings, including horizontal condylar inclination (HCI), immediate mandibular lateral translation (IMLT), progressive mandibular lateral translation (PMLT), top wall, and rear wall, were investigated. Thirty trials were performed for each condylar determinant preset. The standard deviation was calculated for evaluation of reliability and validity. Analysis of variance and Tukey's HSD test were used for statistical analysis ($\alpha = .05$). The reliability readings for the condylar setting at the 10-mm condylotrack distance were most consistent as compared to those at distances of 3 mm and 5 mm. The Denar Mark II had the smallest deviations from the preset values. The authors concluded that the Cadiax Compact is both reliable and valid for the articulators tested.

Chang WSW, Romberg E, Driscoll CF, Tobacco MJ. *J Prosthet Dent* 2004;92:83–89. **References:** 15. **Reprints:** Dr Carl F. Driscoll, Rm 3D08, Baltimore College of Dental Surgery, 666 W Baltimore St, Baltimore, MD 21201. e-mail: cfd001@dental.umaryland.edu—*Eunghwan Kim, Lincoln, NE*